

Apr 15th, 4:35 PM - 5:00 PM

Doxycycline Resistance in Plasmodium falciparum Linked to Single Nucleotide Polymorphisms in the Plasmodium falciparum apicoplast Small Subunit Ribosomal RNA (pfSSrRNA) gene

Amanda Wasko

Dominican University of California

Follow this and additional works at: <http://scholar.dominican.edu/scw>

 Part of the [Biochemistry Commons](#), [Molecular Biology Commons](#), [Parasitology Commons](#), and the [Pathogenic Microbiology Commons](#)

Wasko, Amanda, "Doxycycline Resistance in Plasmodium falciparum Linked to Single Nucleotide Polymorphisms in the Plasmodium falciparum apicoplast Small Subunit Ribosomal RNA (pfSSrRNA) gene" (2016). *Scholarly & Creative Works Conference 2018*. 14. <http://scholar.dominican.edu/scw/scw2016/AllConference/14>

This Event is brought to you for free and open access by The Dominican Experience at Dominican Scholar. It has been accepted for inclusion in Scholarly & Creative Works Conference 2018 by an authorized administrator of Dominican Scholar. For more information, please contact michael.pujals@dominican.edu.

Doxycycline resistance in *Plasmodium falciparum* linked to single nucleotide polymorphisms in the *Plasmodium falciparum* apicoplast small subunit ribosomal RNA (*pfSSrRNA*) gene

Plasmodium falciparum, a protozoan parasite known as malaria, widely impacts human health; thus antimalarial drug investigations are critical. Doxycycline is a commonly used antimalarial prophylactic, but its mechanism of action is unclear. In prokaryotes, doxycycline works as an antibacterial by disrupting protein translation via the small subunit ribosome. Interestingly, *P. falciparum* has a small subunit ribosome of prokaryotic origins in the apicoplast, a plastid-like organelle. Therefore, we hypothesized that doxycycline works in *P. falciparum* by inhibiting protein synthesis via the small subunit ribosomal RNA and that mutations in the gene encoding the *P. falciparum* apicoplast small subunit ribosomal RNA (*pfSSrRNA*) are associated with doxycycline resistance. We generated nine doxycycline-resistant parasite lines with continuous incremental drug pressure over approximately twelve months. Resistant lines had doxycycline IC₅₀ values of ~2.4 μM while the drug-free control had an IC₅₀ of 587 nM, resulting in a ~4 fold decrease in doxycycline susceptibility as measured by 96-hour growth inhibition assays. Sequencing of the *pfssrRNA* gene in a resistant line revealed a g439a single nucleotide polymorphism (SNP). We suspect this mutation reduces parasite susceptibility to doxycycline by altering the molecular structure of the apicoplast SSrRNA, so that doxycycline can no longer inhibit protein synthesis. Future studies involve cloning the resistant lines, sequencing *pfssrRNA* for all resistant lines, and characterizing the drug phenotypes of the resistant lines to doxycycline and a panel of standard antimalarial drugs. Understanding how doxycycline works will help health care professionals better combat doxycycline resistance in the field.